Colorectal Cancer Deaths, by Sex, Maine, 2008

	Number of Deaths	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine Total	270	20.5	18.1 - 23.1	16.3	14.4 - 18.4
Female	125	18.5	15.4 - 22.1	12.7	10.5 - 15.2
Male	145	22.5	19.0 - 26.5	21.3	18.0 - 25.2

Colorectal Cancer Deaths, by Age and Race, Maine, 2004-2008

	Count	Crude Rate		Age-adjusted Rate	
	Count	Rate	95% CI	Rate	95% CI
Age					
00-14 years	NSD	NSD	NSD	NApp	NApp
15-19 years	NSD	NSD	NSD	NApp	NApp
20-24 years	NSD	NSD	NSD	NApp	NApp
25-29 years	NSD	NSD	NSD	NApp	NApp
30-34 years	NSD	NSD	NSD	NApp	NApp
35-39 years	NSD	NSD	NSD	NApp	NApp
40-44 years	19	3.7	2.2 - 5.8	NApp	NApp
45-49 years	40	7.3	5.2 - 9.9	NApp	NApp
50-54 years	62	11.8	9.1 - 15.2	NApp	NApp
55-59 years	93	19.9	16.1 - 24.4	NApp	NApp
60-64 years	120	33.5	27.8 - 40.1	NApp	NApp
65-69 years	129	47.8	39.9 - 56.8	NApp	NApp
70-74 years	161	70.8	60.3 - 82.6	NApp	NApp
80-84 years	216	147.6	128.6 - 168.6	NApp	NApp
85+ years	353	269.2	241.9 - 298.8	NApp	NApp
Race					
White	2,999	23.7	22.8 - 24.5	19.7	19.0 - 20.4
Black	NSD	NSD	NSD	NSD	NSD
Other*	21	10.3	6.4 - 15.7	21.0	12.2 - 33.2

Data Source: Surveillance, Epidemiology, and End Results SEER Program, National Cancer Institute, Underlying mortality data provided by National Center for Health Statistics. Rates are calculated using SEER*Stat Version 7.0.5.

Cancer Deaths: Deaths with malignant cancer as the underlying cause of death. Colorectal Cancer: SEER Cause of Death Recode: 21040-21050 (which include 1999-2009: ICD-10 codes C18, C26.0,C19-C20; 1993-1998: ICD-9 codes 153, 159.0,154.0-154.1).

Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

NSD = Non sufficient data. Data not available due to too few cases.

Napp: Not applicableage-adjusted rates are not calculated for age-specific groups.

^{*}Other Race includes American Indian/AK Native and Asian/Pacific Islander